

Another four critical points theorem

BIAGIO RICCERI

Dedicated to the memory of Ky Fan, with my immense esteem and admiration

Abstract: In this paper, making use of Theorem 2 of [5], we establish a new four critical points theorem which can be regarded as a companion to Theorem 1 of [4]. We also present an application to the Dirichlet problem for a class of quasilinear elliptic equations.

Key words: Critical point, global minimum, multiplicity, minimax inequality, Dirichlet problem.

2010 Mathematics Subject Classification: 47J10, 47J30, 58E05, 49J35, 35J92.

The aim of this paper is to establish a new four critical points theorem (Theorem 1 below) that can be regarded as a companion to Theorem 1 of [4].

As in [4], our key tool is the multiplicity result on global minima established in [3]. The use of such a result requires the validity of a strict minimax inequality which is explicitly assumed in [4].

A novelty of Theorem 1 is that no minimax inequality appears among the hypotheses. This is possible thanks to the use of Theorem 2 of the very recent [5] which just highlights a rather general situation where the strict minimax inequality occurs.

In other words, Theorem 1 should be regarded as the fruit that one obtains by combining the underlying ideas of [4] with Theorem 2 of [5].

For the reader convenience, we start just recalling Theorem 2 of [5].

First, we introduce the following notations.

If X is a non-empty set and $\Gamma, \Psi, \Phi : X \rightarrow \mathbf{R}$ are three given functions, for each $\mu > 0$ and $r \in]\inf_X \Phi, \sup_X \Phi[$, we put

$$\alpha(\mu\Gamma + \Psi, \Phi, r) = \inf_{x \in \Phi^{-1}(]-\infty, r])} \frac{\mu\Gamma(x) + \Psi(x) - \inf_{\Phi^{-1}(]-\infty, r])}(\mu\Gamma + \Psi)}{r - \Phi(x)}$$

and

$$\beta(\mu\Gamma + \Psi, \Phi, r) = \sup_{x \in \Phi^{-1}(]r, +\infty[)} \frac{\mu\Gamma(x) + \Psi(x) - \inf_{\Phi^{-1}(]-\infty, r])}(\mu\Gamma + \Psi)}{r - \Phi(x)}.$$

When $\Psi + \Phi$ is bounded below, for each $r \in]\inf_X \Phi, \sup_X \Phi[$ such that

$$\inf_{x \in \Phi^{-1}(]-\infty, r])} \Gamma(x) < \inf_{x \in \Phi^{-1}(r)} \Gamma(x) ,$$

we put

$$\mu^*(\Gamma, \Psi, \Phi, r) = \inf \left\{ \frac{\Psi(x) - \gamma + r}{\eta_r - \Gamma(x)} : x \in X, \Phi(x) < r, \Gamma(x) < \eta_r \right\} ,$$

where

$$\gamma = \inf_{x \in X} (\Psi(x) + \Phi(x))$$

and

$$\eta_r = \inf_{x \in \Phi^{-1}(r)} \Gamma(x) .$$

THEOREM A ([5], Theorem 2). - *Let X be a topological space and $\Gamma, \Psi, \Phi : X \rightarrow \mathbf{R}$ three sequentially lower semicontinuous functions, with Γ also sequentially inf-compact, satisfying the following conditions:*

- (a) $\inf_{x \in X} (\mu \Gamma(x) + \Psi(x)) = -\infty$ for all $\mu > 0$;
- (b) $\inf_{x \in X} (\Psi(x) + \Phi(x)) > -\infty$;
- (c) *there exists $r \in]\inf_X \Phi, \sup_X \Phi[$ such that*

$$\inf_{x \in \Phi^{-1}(]-\infty, r])} \Gamma(x) < \inf_{x \in \Phi^{-1}(r)} \Gamma(x) .$$

Under such hypotheses, for each $\mu > \max\{0, \mu^*(\Gamma, \Psi, \Phi, r)\}$, one has

$$\alpha(\mu \Gamma + \Psi, \Phi, r) = 0$$

and

$$\beta(\mu \Gamma + \Psi, \Phi, r) > 0 .$$

As we said, the key tool in the proof of Theorem 1 is provided by the following

THEOREM B ([3], Theorem 1). - *Let X be a topological space, $A \subseteq \mathbf{R}$ an open interval and $P : X \times A \rightarrow \mathbf{R}$ a function satisfying the following conditions:*

- (a₁) *for each $x \in X$, the function $P(x, \cdot)$ is quasi-concave and continuous ;*
- (a₂) *for each $\lambda \in A$, the function $P(\cdot, \lambda)$ is lower semicontinuous and inf-compact ;*
- (a₃) *one has*

$$\sup_{\lambda \in A} \inf_{x \in X} P(x, \lambda) < \inf_{x \in X} \sup_{\lambda \in A} P(x, \lambda) .$$

Under such hypotheses, there exists $\lambda^* \in A$ such that the function $P(\cdot, \lambda^*)$ has at least two global minima.

Here is our main result:

THEOREM 1. - Let X be a reflexive real Banach space; $I : X \rightarrow \mathbf{R}$ a sequentially weakly lower semicontinuous and coercive C^1 functional whose derivative admits a continuous inverse on X^* ; $J, \Psi, \Phi : X \rightarrow \mathbf{R}$ three C^1 functionals with compact derivative satisfying the following conditions:

$$\liminf_{\|x\| \rightarrow +\infty} \frac{J(x)}{I(x)} \geq 0, \quad \limsup_{\|x\| \rightarrow +\infty} \frac{J(x)}{I(x)} < +\infty, \quad (1)$$

$$\liminf_{\|x\| \rightarrow +\infty} \frac{\Psi(x)}{I(x)} = -\infty, \quad (2)$$

$$\inf_{x \in X} (\Psi(x) + \lambda \Phi(x)) > -\infty \quad (3)$$

for all $\lambda > 0$. Moreover, assume that there exist a strict local minimum x_0 of I , with $I(x_0) = J(x_0) = \Psi(x_0) = \Phi(x_0) = 0$, and another point $x_1 \in X$ such that

$$\max\{J(x_1), \Psi(x_1), \Phi(x_1)\} < 0, \quad (4)$$

$$\min \left\{ \liminf_{x \rightarrow x_0} \frac{J(x)}{I(x)}, \liminf_{x \rightarrow x_0} \frac{\Phi(x)}{I(x)} \right\} \geq 0 \quad (5)$$

and

$$\liminf_{x \rightarrow x_0} \frac{\Psi(x)}{I(x)} > -\infty.$$

Under such hypotheses, for each ν, μ satisfying

$$\nu > \max \left\{ 0, -\frac{I(x_1)}{J(x_1)} \right\} \quad (6)$$

and

$$\mu > \max \left\{ 0, -\liminf_{x \rightarrow x_0} \frac{\Psi(x)}{I(x)}, \inf_{r > \sup_{M_\nu} \Phi} \mu^*(I + \nu J, \Psi, \Phi, r) \right\}, \quad (7)$$

where M_ν is the set of all global minima of $I + \nu J$, there exists $\lambda^* > 0$ such that the functional $\mu(I + \nu J) + \Psi + \lambda^* \Phi$ has at least four critical points. Precisely, among them, one is x_0 as a strict local, not global minimum and two are global minima.

PROOF. First of all, observe that, since X is reflexive, the functionals J, Ψ, Φ are sequentially weakly continuous, being with compact derivative ([6], Corollary 41.9). Fix ν as in (6). For $x \in X \setminus I^{-1}(0)$, we have

$$I(x) + \nu J(x) = I(x) \left(1 + \nu \frac{J(x)}{I(x)} \right)$$

and so, since I is coercive, in view of (1), it follows that

$$\lim_{\|x\| \rightarrow +\infty} (I(x) + \nu J(x)) = +\infty. \quad (8)$$

By the reflexivity of X again, this implies that the set M_ν is non-empty and bounded. As a consequence, Φ is bounded in M_ν . Also, by (2) and (3), we have $\sup_X \Phi = +\infty$. Now, fix μ as in (7). Let $r > \sup_{M_\nu} \Phi$ be such that $\mu > \mu^*((I + \nu J), \Psi, \Phi, r)$. Since $\Phi^{-1}(r)$ is non-empty and sequentially weakly closed, there exists $\bar{x} \in \Phi^{-1}(r)$ such that

$$I(\bar{x}) + \nu J(\bar{x}) = \inf_{x \in \Phi^{-1}(r)} (I(x) + \nu J(x)) .$$

The choice of r implies that $\bar{x} \notin M_\nu$. So, we infer that

$$\inf_{x \in \Phi^{-1}([-\infty, r])} (I(x) + \nu J(x)) < \inf_{x \in \Phi^{-1}(r)} (I(x) + \nu J(x)) .$$

Moreover, by (2), there exists a sequence $\{x_n\}$ in X such that

$$\lim_{n \rightarrow \infty} \|x_n\| = +\infty , \quad \lim_{n \rightarrow \infty} \frac{\Psi(x_n)}{I(x_n)} = -\infty . \quad (9)$$

For any $\rho \in \mathbf{R}$ and for n large enough, we have

$$\rho(I(x_n) + \nu J(x_n)) + \Psi(x_n) = (I(x_n) + \nu J(x_n)) \left(\rho + \frac{\frac{\Psi(x_n)}{I(x_n)}}{1 + \nu \frac{J(x_n)}{I(x_n)}} \right) . \quad (10)$$

Clearly, from (1), (8), (9) and (10), it follows that

$$\lim_{n \rightarrow \infty} (\mu(I(x_n) + \nu J(x_n)) + \Psi(x_n)) = -\infty .$$

So, if we consider X endowed with the weak topology, all the assumptions of Theorem A (with $\Gamma = I + \nu J$) are satisfied, and so we have

$$\alpha(\mu(I + \nu J) + \Psi, \Phi, r) < \beta(\mu(I + \nu J) + \Psi, \Phi, r) .$$

But, by Theorem 1 of [1], this inequality is equivalent to

$$\sup_{\lambda \geq 0} \inf_{x \in X} ((\mu(I(x) + \nu J(x)) + \Psi(x) + \lambda(\Phi(x) - r))) < \inf_{x \in X} \sup_{\lambda \geq 0} ((\mu(I(x) + \nu J(x)) + \Psi(x) + \lambda(\Phi(x) - r))) .$$

At this point, after observing that, in view of (3) and (8), one has

$$\lim_{\|x\| \rightarrow +\infty} (\mu(I(x) + \nu J(x)) + \Psi(x) + \lambda \Phi(x)) = +\infty \quad (11)$$

for all $\lambda > 0$, we realize that we can apply Theorem B, with $A =]0, +\infty[$, considering X with the weak topology again and taking

$$P(x, \lambda) = \mu(I(x) + \nu J(x)) + \Psi(x) + \lambda(\Phi(x) - r) .$$

Therefore, there exists $\lambda^* > 0$ such that the functional $\mu(I + \nu J) + \Psi + \lambda^* \Phi$ has at least two global minima. Now, choose $\epsilon, \sigma > 0$ so that

$$\liminf_{x \rightarrow x_0} \frac{\Psi(x)}{I(x)} > -\mu + \epsilon$$

and

$$\sigma < \frac{\epsilon}{\mu\nu + \lambda^*} .$$

In view of (5) and recalling that x_0 is a strict local minimum of I (with $I(x_0) = 0$), we can find a neighbourhood V of x_0 such that, for each $x \in V \setminus \{x_0\}$, one has

$$I(x) > 0 ,$$

$$\frac{\Psi(x)}{I(x)} > -\mu + \epsilon ,$$

$$\frac{J(x)}{I(x)} > -\sigma ,$$

and

$$\frac{\Phi(x)}{I(x)} > -\sigma .$$

Consequently, for each $x \in V \setminus \{x_0\}$, we have

$$\begin{aligned} \mu(I(x) + \nu J(x)) + \Psi(x) + \lambda^* \Phi(x) &= I(x) \left(\mu + \mu\nu \frac{J(x)}{I(x)} + \frac{\Psi(x)}{I(x)} + \lambda^* \frac{\Phi(x)}{I(x)} \right) > \\ &> I(x)(\epsilon - \sigma(\mu\nu + \lambda^*)) > 0 . \end{aligned}$$

Hence, since $\mu(I(x_0) + \nu J(x_0)) + \Psi(x_0) + \lambda^* \Phi(x_0) = 0$, it follows that x_0 is a strict local minimum of the functional $\mu(I + \nu J) + \Psi + \lambda^* \Phi$. On the other hand, in view of (4) and (6), we have

$$\mu(I(x_1) + \nu J(x_1)) + \Psi(x_1) + \lambda^* \Phi(x_1) < 0 ,$$

and hence x_0 is not a global minimum of the functional $\mu(I + \nu J) + \Psi + \lambda^* \Phi$. Now, we remark that this functional, due to (11) and to our assumptions on I, J, Φ, Ψ turns out to satisfy the Palais-Smale condition ([6], Example 38.25). Summarizing: the functional $\mu(I + \nu J) + \Psi + \lambda^* \Phi$ is C^1 , satisfies the Palais-Smale condition, has at least two global minima and admits x_0 as a local, not global minimum. At this point, we can invoke Theorem (1.ter) of [2] to ensure the existence of a fourth critical point for the same functional, and the proof is complete. $\triangle$

Now, we are going to present an application of Theorem 1 to quasilinear elliptic equations.

So, let $\Omega \subset \mathbf{R}^n$ be a bounded domain with smooth boundary and let $p > 1$. On the Sobolev space $W_0^{1,p}(\Omega)$, we consider the norm

$$\|u\| = \left(\int_{\Omega} |\nabla u(x)|^p dx \right)^{\frac{1}{p}}.$$

If $n \geq p$, we denote by $\mathcal{A}$ the class of all Carathéodory functions $f : \Omega \times \mathbf{R} \rightarrow \mathbf{R}$ such that

$$\sup_{(x,\xi) \in \Omega \times \mathbf{R}} \frac{|f(x,\xi)|}{1 + |\xi|^q} < +\infty,$$

where $0 < q < \frac{pn-n+p}{n-p}$ if $p < n$ and $0 < q < +\infty$ if $p = n$. While, when $n < p$, we denote by $\mathcal{A}$ the class of all Carathéodory functions $f : \Omega \times \mathbf{R} \rightarrow \mathbf{R}$ such that, for each $r > 0$, the function $x \rightarrow \sup_{|\xi| \leq r} |f(x,\xi)|$ belongs to $L^1(\Omega)$.

Given $f \in \mathcal{A}$, consider the following Dirichlet problem

$$\begin{cases} -\operatorname{div}(|\nabla u|^{p-2} \nabla u) = f(x, u) & \text{in } \Omega \\ u = 0 & \text{on } \partial\Omega. \end{cases} \quad (P_f)$$

Let us recall that a weak solution of (P_f) is any $u \in W_0^{1,p}(\Omega)$ such that

$$\int_{\Omega} |\nabla u(x)|^{p-2} \nabla u(x) \nabla v(x) dx - \int_{\Omega} f(x, u(x)) v(x) dx = 0$$

for all $v \in W_0^{1,p}(\Omega)$.

The functionals $T, J_f : W_0^{1,p}(\Omega) \rightarrow \mathbf{R}$ defined by

$$T(u) = \frac{1}{p} \|u\|^p$$

$$J_f(u) = \int_{\Omega} F(x, u(x)) dx,$$

where

$$F(x, \xi) = \int_0^{\xi} f(x, t) dt,$$

are C^1 with derivatives given by

$$T'(u)(v) = \int_{\Omega} |\nabla u(x)|^{p-2} \nabla u(x) \nabla v(x) dx$$

$$J'_f(u)(v) = \int_{\Omega} f(x, u(x)) v(x) dx$$

for all $u, v \in W_0^{1,p}(\Omega)$. Consequently, the weak solutions of problem (P_f) are exactly the critical points in $W_0^{1,p}(\Omega)$ of the functional $T - J_f$ which is called the energy functional of problem (P_f) . Moreover, J'_f is compact, while T' is a homeomorphism between $W_0^{1,p}(\Omega)$ and its dual.

The announced application of Theorem 1 is as follows:

THEOREM 2. - *Let $q > p$, with $q < \frac{pn}{n-p}$ when $n > p$, and let $f, g, h : \Omega \times \mathbf{R} \rightarrow \mathbf{R}$ be three functions belonging to $\mathcal{A}$ and satisfying the following conditions:*

$$\lim_{\xi \rightarrow +\infty} \frac{\inf_{x \in \Omega} F(x, \xi)}{\xi^p} = +\infty, \quad \limsup_{|\xi| \rightarrow +\infty} \frac{\sup_{x \in \Omega} F(x, \xi)}{|\xi|^q} < +\infty, \quad (12)$$

$$\lim_{|\xi| \rightarrow +\infty} \frac{\inf_{x \in \Omega} G(x, \xi)}{|\xi|^q} = +\infty, \quad (13)$$

$$\limsup_{|\xi| \rightarrow +\infty} \frac{\sup_{x \in \Omega} H(x, \xi)}{|\xi|^p} \leq 0, \quad \liminf_{|\xi| \rightarrow +\infty} \frac{\inf_{x \in \Omega} H(x, \xi)}{|\xi|^p} > -\infty, \quad (14)$$

$$\limsup_{\xi \rightarrow 0} \frac{\sup_{x \in \Omega} F(x, \xi)}{|\xi|^p} < +\infty, \quad (15)$$

$$\liminf_{\xi \rightarrow 0} \frac{\inf_{x \in \Omega} G(x, \xi)}{|\xi|^p} \geq 0, \quad (16)$$

$$\limsup_{\xi \rightarrow 0} \frac{\sup_{x \in \Omega} H(x, \xi)}{|\xi|^p} \leq 0. \quad (17)$$

Finally, assume that there exist a measurable set $B \subset \Omega$, with $\text{meas}(B) > 0$, and $\xi_1 \in \mathbf{R}$ such that

$$\max\{-F(x, \xi_1), G(x, \xi_1), -H(x, \xi_1)\} < 0$$

for all $x \in B$.

Under such hypotheses, for each $\nu > 0$ large enough, there exists $\epsilon_\nu > 0$ with the following property: for each $\epsilon \in]0, \epsilon_\nu[$ there exists $\lambda^* > 0$ such that the problem

$$\begin{cases} -\text{div}(|\nabla u|^{p-2} \nabla u) = \epsilon f(x, u) - \lambda^* g(x, u) + \nu h(x, u) & \text{in } \Omega \\ u = 0 & \text{on } \partial\Omega \end{cases}$$

has at least three non-zero weak solutions, two of which are global minima in $W_0^{1,p}(\Omega)$ of the corresponding energy functional.

PROOF. First, observe that from the first assumption in (12) it follows

$$\limsup_{\|u\| \rightarrow +\infty} \frac{J_f(u)}{\|u\|^p} = +\infty.$$

This is proved in the proof of Theorem 4 of [5], and so we do not repeat the argument here. Moreover, from the second assumption in (12) and from (13), it clearly follows that,

for each $\lambda > 0$, the function $\lambda G - F$ is bounded below in $\mathbf{R}$ (see [5] again), and so the functional $\lambda J_g - J_f$ is bounded below in $W_0^{1,p}(\Omega)$. Moreover, by (14), there is $c > 0$ and, for each $\epsilon > 0$, another $c_\epsilon > 0$, such that

$$-c(|\xi|^p + 1) \leq H(x, \xi) \leq \epsilon|\xi|^p + c_\epsilon$$

for all $(x, \xi) \in \Omega \times \mathbf{R}$. This clearly implies that

$$\limsup_{\|u\| \rightarrow +\infty} \frac{J_h(u)}{\|u\|^p} \leq 0$$

and

$$\liminf_{\|u\| \rightarrow +\infty} \frac{J_h(u)}{\|u\|^p} > -\infty .$$

Furthermore, by (15) and by the second assumption in (12), there is a constant $c_1 > 0$ such that

$$F(x, \xi) \leq c_1(|\xi|^p + |\xi|^q)$$

for all $(x, \xi) \in \Omega \times \mathbf{R}$. Since $q > p$, this implies that

$$\limsup_{u \rightarrow 0} \frac{J_f(u)}{\|u\|^p} < +\infty .$$

Now, suppose $n \geq p$. By (16) and (17), taking into account that $g, h \in \mathcal{A}$, for some $r > p$ and for each $\epsilon > 0$ there is $d_\epsilon > 0$ such that

$$G(x, \xi) \geq -\epsilon|\xi|^p - d_\epsilon|\xi|^r$$

and

$$H(x, \xi) \leq \epsilon|\xi|^p + d_\epsilon|\xi|^r$$

for all $(x, \xi) \in \Omega \times \mathbf{R}$. From this, we get

$$\liminf_{u \rightarrow 0} \frac{J_g(u)}{\|u\|^p} \geq 0$$

and

$$\limsup_{u \rightarrow 0} \frac{J_h(u)}{\|u\|^p} \leq 0 .$$

In the case $n < p$, we get again these two inequalities thanks to (16) and (17) and to the continuous embedding of $W_0^{1,p}(\Omega)$ into $C^0(\overline{\Omega})$. Now, let $\omega \in L^1(\Omega)$ be such that

$$\max\{|F(x, \xi)|, |G(x, \xi)|, |H(x, \xi)|\} \leq \omega(x)$$

for all $(x, \xi) \in \Omega \times [-|\xi_1|, |\xi_1|]$. Next, choose a closed set $C \subset B$, an open set $D \subset \Omega$, with $C \subset D$, and $\theta \in \mathbf{R}$ in such a way that

$$\eta := \max \left\{ - \int_C F(x, \xi_1) dx, \int_C G(x, \xi_1) dx, - \int_C H(x, \xi_1) dx \right\} < \theta < 0$$

and

$$\int_{D \setminus C} \omega(x) dx < \theta - \eta .$$

Finally, let $v_1 : \Omega \rightarrow [-|\xi_1|, |\xi_1|]$ be a function belonging to $W_0^{1,p}(\Omega)$ such that $v_1(x) = \xi_1$ for all $x \in C$ and $v_1(x) = 0$ for all $x \in \Omega \setminus D$. Clearly, we have

$$\max \left\{ - \int_{\Omega} F(x, v_1(x)) dx, \int_{\Omega} F(x, v_1(x)) dx, - \int_{\Omega} H(x, v_1(x)) dx \right\} < \eta + (\theta - \eta) < 0 .$$

At this point, the conclusion comes directly from that of Theorem 1 applied taking $I(u) = \frac{1}{p} \|u\|^p$, $J(u) = -J_h(u)$, $\Psi(u) = -J_f(u)$, $\Phi(u) = J_g(u)$ for all $u \in W_0^{1,p}(\Omega)$. $\triangle$

References

- [1] G. CORDARO, *On a minimax problem of Ricceri*, J. Inequal. Appl., **6** (2001), 261-285.
- [2] N. GHOUSSEUB and D. PREISS, *A general mountain pass principle for locating and classifying critical points*, Ann. Inst. H. Poincaré Anal. Non Linéaire, **6** (1989), 321-330.
- [3] B. RICCERI, *Multiplicity of global minima for parametrized functions*, Rend. Lincei Mat. Appl., **21** (2010), 47-57.
- [4] B. RICCERI, *A class of nonlinear eigenvalue problems with four solutions*, J. Nonlinear Convex Anal., **11** (2010), 503-511.
- [5] B. RICCERI, *A further refinement of a three critical points theorem*, Nonlinear Anal., **74** (2011), 7446-7454.
- [6] E. ZEIDLER, *Nonlinear functional analysis and its applications*, vol. III, Springer-Verlag, 1985.

Department of Mathematics
University of Catania
Viale A. Doria 6
95125 Catania
Italy
e-mail address: ricceri@dmf.unict.it